

Draft Genome Sequence of *Microvirga* sp. Strain BSC39, Isolated from Biological Soil Crust of Moab, Utah

Alexis C. Bailey,^a Matthew Kellom,^a Amisha T. Poret-Peterson,^a Kathryn Noonan,^a Hilairy E. Hartnett,^{a,b} Jason Raymond^a

School of Earth and Space Exploration, Arizona State University, Tempe, Arizona, USA^a; Department of Chemistry and Biochemistry, Arizona State University, Tempe, Arizona, USA^b

A.C.B. and M.K. contributed equally to this work.

***Microvirga* sp. BSC39 was isolated from a biological soil crust near Moab, Utah. The strain appears to be capable of chemotaxis and exopolysaccharide synthesis for biofilm adhesion. The BSC39 genome contains iron siderophore uptake and hydrolysis enzymes; however, it lacks siderophore synthesis pathways, suggesting the uptake of siderophores produced by neighboring microbes.**

Received 7 October 2014 Accepted 8 October 2014 Published 13 November 2014

Citation Bailey AC, Kellom M, Poret-Peterson AT, Noonan K, Hartnett HE, Raymond J. 2014. Draft genome sequence of *Microvirga* sp. strain BSC39, isolated from biological soil crust of Moab, Utah. *Genome Announc.* 2(6):e01197-14. doi:10.1128/genomeA.01197-14.

Copyright © 2014 Bailey et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](https://creativecommons.org/licenses/by/3.0/).

Address correspondence to Jason Raymond, jason.raymond@asu.edu.

Microvirga sp. strain BSC39 was collected in an environmental sample from the Green Butte site near Moab, Utah (N38°42'56.2", W109°41'32") in May 2009 from a biological soil crust (BSC) (1). BSCs are microbial communities that form in arid and semiarid environments and supply bioavailable carbon (2), nitrogen (3), and metal-containing compounds (4) to their surroundings via carbon and nitrogen fixation as well as siderophore synthesis (5; K. Noonan, A. T. Poret-Peterson, A. D. Potrafka, A. D. Anbar, F. Garcia-Pichel, and H. E. Hartnett, unpublished data). Here, we present the draft genome of *Microvirga* sp. BSC39, a heterotrophic, siderophore-hydrolyzing member of *Alphaproteobacteria*.

BSC39 was cultured and isolated on BG-11 agar plates and modified to detect the production of siderophores (6). DNA was extracted from the isolate using the Ultra-Clean Soil DNA Extraction kit, prepped for sequencing using the Illumina TruSeq DNA HT sample prep kit, and sequenced on 21 June 2014 at the Genomics Core of The Biodesign Institute at Arizona State University using the Illumina MiSeq (Illumina RTA 1/18/54). A total of 833,789 300-bp paired-end reads were generated, resulting in 335.1 Mb of raw sequence data. We assembled 18,135 Rho N_{50} unitigs averaging 419 bp in length and remaining reads into 80 gapless scaffolds using the Celera Whole-Genome Shotgun Assembler version 8.1 for a total genome length of 5,655,736 bp (50× coverage) (7). The scaffolds were screened for DNA contaminants and annotated by the NCBI Prokaryotic Genome Annotation Pipeline, which identified 4,793 genes (8). 16S ribosomal analysis showed 99% identity to the 16S sequence of *Microvirga aeralata* from the NCBI 16S Ribosomal RNA Sequences Database using BLAST (9). The G+C content was 62.94%, which is consistent with the *Microvirga* genus (10).

Based on the genome of BSC39, it appears to be an opportunistic heterotrophic member of the soil crust microbial community with a complete TCA cycle and terminal cytochrome oxidases, metabolizing carbon and also nitrogen biomolecules

produced by other BSC microbes. BSC39 has genes for chemotaxis, flagellar motility, and exopolysaccharide biofilm synthesis, which facilitate movement toward and adhesion to areas of favorable nutrient conditions. BSC39 also appears to be capable of iron acquisition via siderophore uptake by iron siderophore ABC transporter permease and siderophore hydrolysis. The genes producing siderophores do not appear to be present within the BSC39 genome, and therefore any siderophores taken up and hydrolyzed for iron acquisition must have been produced by other microbes within the microbial community.

Nucleotide sequence accession numbers. This whole-genome shotgun project has been deposited at DDBJ/EMBL/GenBank under accession number [JPUG00000000](https://www.ncbi.nlm.nih.gov/nuccore/JPUG00000000). The version described in this paper is version JPUG01000000.

ACKNOWLEDGMENTS

We acknowledge support from the NASA Exobiology and Evolutionary Biology Program (NNX08AP61G), NASA Astrobiology Institute at ASU "Follow the Elements" (08-NAI5-0018), and NSF (EAR-0525569).

REFERENCES

- Allen MM, Stanier RY. 1968. Selective isolation of blue-green algae from water and soil. *J. Gen. Microbiol.* 51:203–209. <http://dx.doi.org/10.1099/00221287-51-2-203>.
- Beymer RJ, Klopatek JM. 1991. Potential contribution of carbon by microphytic crusts in pinyon-juniper woodlands. *Arid Soil Res. Rehabil.* 5:187–198. <http://dx.doi.org/10.1080/15324989109381279>.
- Belnap J. 2002. Nitrogen fixation in biological soil crusts from southeast Utah, USA. *Biol. Fertil. Soils* 35:128–135. <http://dx.doi.org/10.1007/s00374-002-0452-x>.
- Beraldi-Campesi H, Hartnett HE, Anbar A, Gordon GW, Garcia-Pichel F. 2009. Effect of biological soil crusts on soil elemental concentrations: implications for biogeochemistry and as traceable biosignatures of ancient life on land. *Geobiology* 7:348–359. <http://dx.doi.org/10.1111/j.1472-4669.2009.00204.x>.
- Noonan K, Anbar AD, Garcia-Pichel F, Poret-Peterson AT, Hartnett HE. 2011. Six siderophore-producing microorganisms identified in biological soil crusts. Abstr. B32A–04. American Geophysical Union, Fall Meeting 2011. AGU, San Francisco, CA.

6. Strauss SL, Day TA, Garcia-Pichel F. 2012. Nitrogen cycling in desert biological soil crusts across biogeographic regions in the Southwestern United States. *Biogeochemistry* 108:171–182. <http://dx.doi.org/10.1007/s10533-011-9587-x>.
7. Myers EW, Sutton GG, Delcher AL, Dew IM, Fasulo DP, Flanigan MJ, Kravitz SA, Mobarry CM, Reinert KH, Remington KA, Anson EL, Bolanos RA, Chou HH, Jordan CM, Halpern AL, Lonardi S, Beasley EM, Brandon RC, Chen L, Dunn PJ, Lai Z, Liang Y, Nusskern DR, Zhan M, Zhang Q, Zheng X, Rubin GM, Adams MD, Venter JC. 2000. A whole-genome assembly of *Drosophila*. *Science* 287:2196–2204. <http://dx.doi.org/10.1126/science.287.5461.2196>.
8. Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Ciufu S, Li W. 2013. Prokaryotic genome annotation pipeline. *In* Beck J, Benson D, Coleman J, Hoepfner M, Johnson M, Maglott M, Mizrahi I, Morris R, Ostell J, Pruitt K, Rubinstein W, Sayers E, Sirotkin K, Tatusova T (ed), *The NCBI handbook*, 2nd ed. National Center for Biotechnology Information, Bethesda, MD. <http://www.ncbi.nlm.nih.gov/books/NBK174280/>.
9. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment search tool. *J. Mol. Biol.* 215:403–410. [http://dx.doi.org/10.1016/S0022-2836\(05\)80360-2](http://dx.doi.org/10.1016/S0022-2836(05)80360-2).
10. Weon HY, Kwon SW, Son JA, Jo EH, Kim SJ, Kim YS, Kim BY, Ka JO. 2010. Description of *Microvirga aerophila* sp. nov. and *Microvirga aerilata* sp. nov., isolated from air, reclassification of *Balneimonas flocculans* Takeda *et al.* 2004 as *Microvirga flocculans* comb. nov. and emended description of the genus *Microvirga*. *Int. J. Syst. Evol. Microbiol.* 60: 2596–2600. <http://dx.doi.org/10.1099/ijs.0.018770-0>.